



# Laser Innovations for Research and Applications

journal homepage: <https://lira.journals.ekb.eg/>

ISSN: Print 3009-6359 / Online 3009-6472



## Antibacterial Effects of Laser in Treating Contaminated Wounds

Esraa Ahmed<sup>1</sup>, Ahmed O. El-Gendy<sup>1,2</sup>, Tarek Mohamed<sup>1\*</sup>

<sup>1</sup>Laser Institute for Research and Applications (LIRA), Beni-Suef University, Beni-Suef 62511, Egypt

<sup>2</sup>Faculty of Pharmacy, Department of Microbiology and Immunology, Beni-Suef University, Beni-Suef 62514, Egypt

### Abstract

Bacterial infections persist as a major worldwide health issue. Regarded as the most distressing factor affecting the wound healing process, resulting in prolonged healing time, increased patient trauma, elevated treatment expenses, and substantial mortality and morbidity. The prevalence of antibiotic resistance among bacteria has led to the emergence of bacterial wound infection as an alarming and complex health issue. The effectiveness of traditional antibiotic treatments is progressively limited, prompting the exploration of alternative therapeutic approaches. Antibacterial Photodynamic Therapy (aPDT) using laser technology has shown great potential in treating persistent bacterial infections. This review aims to provide an extensive overview of different wound treatment strategies and infection management modalities within and surrounding the wound bed as well as the use of aPDT against different bacterial pathogens, with a particular focus on its potential to tackle the worldwide problem of bacterial infections.

**Keywords:** Chronic Wounds; Femtosecond laser; Antibacterial; Photodynamic therapy; Bacterial infection

\*Corresponding author at: Laser Institute for Research and Applications LIRA, Beni-Suef University, Beni-Suef 62511, Egypt

E-mail addresses: tarek\_mohamed1969@lira.bsu.edu.eg

### 1. Introduction

The skin is the largest sensory organ, with a surface area of about 2 square meters (Chang et al., 2020). It's a layered structure made up of cells, fibers, and other components. Skin integrity is difficult to maintain, but it is an essential protection measurement against potentially harmful toxic substances and invading pathogenic microorganisms. Any skin injury provides a warm, moist, and nutritious environment for pathogenic microorganisms to infiltrate and thrive in. Therefore, rapid repair of these injuries is critical to minimize the risk of infection. There are different types of injuries, ranging from minor wounds to wounds that may lead to major disability or even death (Chang et al., 2020).

Wound healing is a distinct biological process characterized by a well-organized sequence of events. The process of healing is crucial in order to reduce tissue damage and ensure sufficient tissue perfusion, oxygenation, proper nourishment, and a moist environment to restore both the anatomical continuity and functionality of the injured skin area (Pierce & Mustoe, 1995). Appropriate wound healing typically progresses through four phases: (i) Hemostasis, (ii) Inflammation, (iii) Proliferation, and (iv) Remodeling of the scar tissue (Schultz et al., 2003; Singer & Clark, 1999).

The process of homeostasis is initiated immediately following an injury by the constriction of blood vessels and the subsequent creation of a fibrin clot (Barchitta et al., 2019). This mechanism effectively stops excessive bleeding and promotes the entry of platelets and growth factors into the body. Inflammation follows, characterized by the invasion of neutrophils to cleanse the wound and the activation of macrophages to facilitate tissue regeneration. Proliferation includes the processes of cell proliferation and migration, which ultimately result in the creation of granulation tissue and the contraction of wounds (Xue & Jackson, 2015). Ultimately, the process of remodeling, which takes many months to years, involves collagen remodeling and the production of scars, augmenting the strength of wounds. Healing results are influenced by a range of factors, including the size of the wound, the quality of the tissue, and the presence of necrosis (Nourian Dehkordi et al., 2019). Inadequate healing or delayed healing might result in the formation of acute or chronic wounds (RASHIDI et al., 2015).

## **2. Types of wounds**

### **2.1. Acute wounds**

Acute wounds are temporary tissue injuries that fully heal within the anticipated timeframe, typically ranging from 8 to 12 weeks. These wounds undergo a systematic progression of wound healing, resulting in minimum scarring and infrequent consequences. There are many types of acute wounds including minor burns, traumatic and surgical wounds (Boateng et al., 2008).

### **2.2. Chronic wounds**

Non-healing or chronic wounds refer to tissue injuries that do not undergo the typical healing process and are frequently trapped in one or more stages of wound healing for an extended duration (usually the inflammatory and/or the proliferative phases). These wounds usually have a prolonged healing time, beyond 12 weeks, and often reoccur. Chronic wounds have a huge socioeconomic burden as a public health concern with an estimated prevalence of 1-2% of the population in developed countries. Patients with chronic wounds suffer from severe emotional and physical trauma. Chronic wounds commonly include various forms, including pressure ulcers, leg ulcers, diabetic ulcers, and nonhealing surgical wounds. Usually, many complications can be associated with chronic wounds such as infection, sepsis, amputation, and death (Boateng et al., 2008).

Repeated trauma, poor primary treatment, underlying physiological conditions, and many Additional factors contribute to the formation of chronic wounds. Therefore, identifying these factors is a primary clinical concern to provide appropriate treatment and prevention modalities (Boateng et al., 2008).

## **3. Factors affecting wound healing**

Many factors may interrupt the healing process, causing prolonged trauma, higher treatment costs as well as significant mortality and morbidity. These factors are broadly classified into: Drugs: Some chemical agents have adverse effects on the healing process, such as Aspirin, Indomethacin, Tolmetin, Cytotoxic agents, radiation, immunosuppressants as well as chemotherapeutic drugs utilized in cancer treatment (Shpichka et al., 2019; Yildirimer et al., 2012).

Systemic factors: such as age (Enoch & Leaper, 2008), malnutrition (Demidova-Rice et al., 2012), as well as some underlying physiological conditions, for example, obesity, arterial disease, or diabetes (Boateng et al., 2008). Local factors: such as wound environment, moisture, local supply of blood and oxygen, as well as Infection; which is the single most crucial inevitable impairment to wound healing (Demidova-Rice et al., 2012; Shpichka et al., 2019).

Despite the presence of bacteria in the skin microbiome and wounds, a critical threshold of bacteria and the formation of clusters of these bacteria known as biofilms can potentially hinder the process of wound healing (Bowler et al., 2001). Infection can lead not only to chronic wounds but also to gangrene, amputation, and even death. The susceptibility to infection is present in both acute and chronic wounds (Gallo & Hooper, 2012). Chronic wounds are even more susceptible to bacterial infection and microbial colonization (Gallo & Hooper, 2012). In chronic wounds, the prolonged skin rupture and slower regenerative process (Nesi-Reis et al., 2018) increase the chances of microbial penetration and consequent proliferation (Lambrechts et al., 2005). Understanding these elements and their impact on wound healing is crucial in developing more effective therapeutic alternatives for chronic wounds infected by bacteria.

### **3.1. Bacterially infected wounds**

Bacterial pathogens create a bioburden on the wound while competing for the limited oxygen and nutrient supply, resulting in weakened tissue that can easily be ruptured (Agren et al., 2000). The unlimited development of bacterial pathogens negatively impacts the progression of wound healing and causes local tissue damage (Frykberg & Banks, 2015; Schultz et al., 2003). This bacterial development starts as contamination with the presence and attachment of bacteria to the wound surface and progresses from acute to chronic colonization where the bacterial pathogen replicates till the wound reaches a critical colonization state and the healing has been impaired (Guo & Dipietro, 2010), to end as an infection with the bacterial pathogens invading the tissue and triggering a systematic response in a quick time course (hours to days) (Bowler et al., 2001; Coggan & Wolfgang, 2012).

A multitude of organisms can be identified within chronic wounds (Dowd et al., 2008; Sprockett et al., 2015). Despite the topical antibiotic treatment, many bacterial pathogens are difficult to manage, especially with the increase in antibiotic-resistant bacterial strains (Boucher et al., 2009; M. E. Jones et al., 2003; R. N. Jones, 2001; Livermore, 2007; Paterson, 2006; Rice, 2006; Spellberg et al., 2008). *Staphylococcus aureus* (*S. aureus*), and *Pseudomonas aeruginosa* (*P. aeruginosa*) are widely recognized as prominent resistant bacteria infecting chronic wounds. (Barsoumian et al., 2015). In the initial phases of the chronic wound, *S. aureus* remains within the superficial wound, while in advanced phases, *P. aeruginosa* is mostly present deeper in the wound (Cardona & Wilson, 2015). have acquired resistance to nearly all antibiotics and disinfectants licensed for clinical use. They are considered a real challenge because of their rapid colonization and survival ability (Barsoumian et al., 2015).

#### **4. Wound Management**

Many wound management modalities have been developed to optimize and accelerate the healing process of chronic wounds (Mustoe et al., 2006). These modalities consider general principles such as: 1) Wound cleaning and appropriate moisturization using non-irritating, nontoxic solutions to minimize any additional trauma or cytotoxicity, 2) Restoring the quality of the wound edge, 3) The elimination of any accumulated hindrances to wound healing such as necrotic tissue and excessive bacterial burden while preserving the vital surrounding tissue (AYELLO et al., 2004). Some of the most known wound healing therapeutic modalities are listed below in Table 1.

**Table 1:** Different wound healing therapeutic modalities:

Management modality	Based on	Reported for treating	Ref.
Systemic administration of drugs	<p>This modality attempts to achieve local therapeutic effects and improve symptom management. Many drugs have been used, such as antibiotics, peptides (anti-tumor necrosis factor <math>\alpha</math> (anti-TNF- <math>\alpha</math>), or <math>\alpha</math>-melanocyte-stimulating hormone), amino acids or their derivatives (proline or N-acetyl cysteine), corticosteroids, aspirin, warfarin, phenytoin, and many growth factors such as: epidermal growth factor (EGF), basic fibroblast growth factor (bFGF), human granulocyte macrophage colony stimulating factor (GMCSF), platelet-derived growth factor (PDGF), or transforming growth factor beta (TGF-<math>\beta</math>).</p> <p>The systemic administration of these drugs has positively affected wound healing even better than their local administration. However, its effectiveness is constrained by the challenges associated with tissue targeting and the off-target adverse effects.</p>	Wide variety of chronic wounds	(Las Heras et al., 2020; Otero-Viñas & Falanga, 2016)
Natural compounds	<p>Several herbal extracts and natural substances, including rosemary oil, curcumin, berberine, Aloe Vera, thyme extract, and honey, have demonstrated antibacterial, angiogenic, and regenerative properties.</p>	Wide variety of chronic wounds	(Hosseinkhani et al., 2017; Nagori & Solanki, 2011; Segev-Zarko et al., 2015; Shrestha et al., 2014)

Management modality	Based on	Reported for treating	Ref.
Debridement strategies	This modality represents a conventional approach to wound management, wherein the elimination of diseased tissue and foreign substances from the wound is achieved through various methods such as surgical, bio-surgical, mechanical, chemical, or enzymatic means. The technique of sharp debridement is widely recognized for its rapidity and efficacy.	Pressure, diabetes, and venous related ulceration (VLU)	(Steed, 2004)
Compression therapy	This modality helps to reduce ulceration by applying external and gradual pressure with special compression bandages systems. It is imperative to conduct a thorough clinical assessment prior to implementing any type of compression therapy, as it has the potential to result in severe consequences, such as limb gangrene.	Lower limb ulcers	(Cullum et al., 2001)
Skin grafting	<p>This modality relies on using Composite tissue and artificial skin dressings to sustain wound closure.</p> <p>This procedure aims to reconstruct the skin tissue, especially in cases of thermal and burn injuries, providing sufficient wound covering and protection against bacteria.</p>	Venous, neuropathic and pressure ulcers	(Las Heras et al., 2020)

## Laser Innovations for Research and Applications

Management modality	Based on	Reported for treating	Ref.
Dressing therapies	<p>This approach employs biomaterial-based systems derived from biological sources, including fibrin, gelatin, collagen, chitosan, starch, and lactose, as well as synthetic chemicals like polyglycolic acid (PGA) and polylactic acid (PLA) as wound dressings.</p> <p>Dressing therapies serve to facilitate the regeneration of new tissue while simultaneously reducing pain, delivering moisturization, preventing friction or shear, and safeguarding both the peri-ulcer tissue and the skin. When selecting a suitable wound dressing, it is important to take into account the present stage of wound healing. Numerous contemporary dressings integrate various elements of wound bed preparation.</p>	Wide variety of chronic and burn wounds	(Las Heras et al., 2020)
Negative pressure wound therapy	<p>Another non-invasive traditional modality, that works by the removal of wound exudates with a vacuum device and generally is not used alone but applied after debridement.</p> <p>This modality is helpful in reducing edema, increasing local perfusion, promoting angiogenesis, decreasing overall wound size, and enhancing granulation tissue formation.</p>	used to treat a wide range of chronic wounds	(Glass et al., 2014; Lalezari et al., 2017)
Electrical stimulation	<p>Electromagnetic pulses are utilized as an adjuvant method to augment the treatment of chronic wounds. Research has demonstrated that it has the capacity to augment fibroblast activation and bolster collagen synthesis.</p>	Various types of chronic wounds	(Khouri et al., 2017; Kloth, 2014)



Management modality	Based on	Reported for treating	Ref.
Hyperbaric oxygen therapy	This modality involves the administration of medication by the elevation of atmospheric pressure within a chamber, while the patient is undergoing 100% oxygenation. This approach has the potential to enhance neovascularization, minimize proinflammatory enzymes, and stimulate the synthesis of collagen and growth factors. Hyperbaric oxygen has been associated with adverse outcomes such as seizures and pneumothorax.	Diabetic and vascular insufficiency ulcers	(Thom, 2009)
Shockwaves	Shockwaves modalities promote angiogenesis and reduce inflammation.	VLU	(Cooper & Bachoo, 2018; Qureshi et al., 2011)
Radiant heat dressing	This modality uses heat to improve oxygen flow, tissue oxygenation and increases subcutaneous oxygen tension.	Mainly postoperative wounds, diabetic, and pressure ulcers	(Price et al., 2000)

## Laser Innovations for Research and Applications

Management modality	Based on	Reported for treating	Ref.
Ultrasound therapy	<p>This modality relies on the mechanical anti-inflammatory effect of both high-frequency and low-frequency ultrasounds, as they reduce macrophages.</p>	Pressure ulcers, diabetic ulcers, and VLU	(Kavros & Coronado, 2018)
Hydrotherapy	<p>Hydrotherapy is a unique wound treatment modality that provides an optimal healing environment. This modality is a form of mechanical debridement technique that aims to clear the wound from any cellular debris.</p>	Pressure, VLC and other chronic wounds.	(Atkin & Ousey, 2016)
Electromagnetic therapy	<p>This modality works by promoting cytokine synthesis.</p>	Ischemic, pressure, and VLU	(Kwan et al., 2015)
Low Level Laser Therapy, photo-biostimulation, or Photo-biomodulation	<p>Light emitted by light-emitting diodes (LEDs), lasers, and other light sources across the wavelength range spanning from vision to infrared has been used as a wound-healing modality.</p> <p>This modality can stimulate various biological processes with the least systemic side effects such as: 1) Increase blood perfusion, and vasodilation, 2) Reduce inflammation and neutrophil infiltration, 3) Enhance fibroblast proliferation, wound reepithelization, and closure, 4) Elevate ATP synthesis, oxygen generation, and protein synthesis, as well as 5) Induce mitotic cellular proliferation.</p>	Wide variety of acute and chronic wounds	(Amini et al., 2021; Freidouni et al., 2015; Gao & Xing, 2009; Kuffler, 2016; Mizutani et al., 2004; RASHIDI et al., 2015; Sefati et al., 2018)

#### **4.1. Infection management within and surrounding the wound bed**

Any wound can develop into a chronic wound even in immune-competent individuals. Thus, appropriate wound treatment modalities aim to eliminate the causing factors of chronic wound development that hinder the healing process. These factors include ischemia and bacterial infection (AYELLO et al., 2004).

Despite being a part of the skin's microbiota, the continuous proliferation of bacterial pathogens leads to the accumulation of a bioburden on the wound (Bowler et al., 2001). Bacterial burden over  $10^5$ - $10^6$  bacteria colony forming units per gram of tissue, or when any amount of  $\beta$ -hemolytic streptococci is present (Sibbald et al., 2000). Hence, it is imperative to decrease the amount of bacteria present in a wound in order to decrease the levels of local and systemic inflammatory substances, hence managing the extended inflammatory stage of chronic wounds (Sibbald et al., 2000). In chronic wounds, topical antibiotics and surgical debridement have been observed to reduce bacterial populations effectively (AYELLO et al., 2004).

##### **4.1.1. Antibiotics**

Antibiotics have saved countless lives since their discovery. Some antibiotics are synthetic, while others are formed naturally as some microbes produce substances that inhibit or kill other microbes competing for the same resources. These products, if successfully extracted and mass-produced, can be used as natural antibiotics drugs or modified as semi-synthetic antibiotics. They remain an essential tool for treating and controlling infectious diseases. Various bacteriostatic or bactericidal antibiotic classes such as quinolones, tetracyclines, aminoglycosides, and cephalosporins can assist wound healing (Kohanski et al., 2010).

The prevalence of antibiotic resistance among bacteria has led to the emergence of bacterial wound infection as a significant and complex health issue (Rai et al., 2016). Although many bacteria are still susceptible to antibiotics (Rice, 2008), about 70% of wound-infecting bacteria are resistant to a minimum of one antibiotic (Friedman et al., 2016). *S. aureus* and *P. aeruginosa* are among these bacteria (Rice, 2008). Infections caused by antibiotic-resistant bacteria are exceptionally hard to treat. These infections are usually associated with a prolonged hospital stay and an approximately two-fold higher death rate (Fang

et al., 2016), with a predicted mortality of 10 million people per year by 2050 (Fang et al., 2016). Since the rate of acquisition of resistance is faster than the clinical introduction of new antibiotics, the development of alternative antimicrobial techniques to treat these infections is becoming necessary for public health throughout the world (Jackson et al., 2018; Kmietowicz, 2017; Kraus, 2008). Amidst the ongoing antibiotic crisis, several advancements in biotechnology, genetic engineering, and synthetic chemistry have introduced novel alternatives to traditional antibiotics for wound management, some are listed below in Table 2.

**4.1.2. Alternative antibacterial therapeutic strategies**

The following antibacterial therapeutic alternatives are arranged in three categories: (1) Naturally existing alternatives, (2) Synthetically generated strategies, and (3) Biotechnology-based strategies.

*Table 2 Different alternative antibacterial therapeutic strategies.*

<b>(1) Naturally Existing Alternatives strategies</b>					
Strategy	Based on				
Phage therapy	<p>Bacteriophages (phages) are viruses that propagate at the expense of bacteria. Their use as an antimicrobial agent is much older than the antibiotic (Gravitz, 2012).</p> <p>Bacteriophages adhere to the bacterial cell wall and introduce their genetic material into the cytoplasm, thereby taking control of the bacterial cell machinery. This process facilitates the synthesis of phage components and the generation of novel phages within the bacterial infection. Over time, bacterial cells undergo dissolution, subsequently leading to the release of phage progeny, which can initiate a subsequent infection cycle (Abedon et al., 2011).</p>				
	<table border="1" style="width: 100%;"> <tr> <th style="width: 50%;">Advantages over antibiotics</th> <th style="width: 50%;">Possible disadvantages</th> </tr> <tr> <td>                     Self-replicating pharmaceuticals (Labrie et al., 2010).                      Bacteriophages can distinguish between different bacterial populations, thus                 </td> <td>                     High specificity.                      Bacterial pathogens can develop resistance to phages (Abedon et al., 2011).                 </td> </tr> </table>	Advantages over antibiotics	Possible disadvantages	Self-replicating pharmaceuticals (Labrie et al., 2010). Bacteriophages can distinguish between different bacterial populations, thus	High specificity. Bacterial pathogens can develop resistance to phages (Abedon et al., 2011).
	Advantages over antibiotics	Possible disadvantages			
Self-replicating pharmaceuticals (Labrie et al., 2010). Bacteriophages can distinguish between different bacterial populations, thus	High specificity. Bacterial pathogens can develop resistance to phages (Abedon et al., 2011).				

	<p>provide an alternative yet selective antimicrobial strategy (Abedon, 2015; J. R. Clark &amp; March, 2006). Responsive to genetic engineering, thus many of its limitations have been overcome with genetic engineering enabling their use in innovative ways (Braff et al., 2016).</p>	<p>High toxicity with the accumulation of endotoxins and pyrogenic substances after bacterial lysis. Their pharmacokinetics, challenges in phage preparation, its stability, and immunogenicity (Labrie et al., 2010).</p>
Strategy	Based on	
Antimicrobial peptides (AMPs)	As the initial line of defense, host-defense peptides (HDPs) and antimicrobial peptides (AMPs) are released as soon as the pathogen invades the body (Brogden, 2005; Hancock & Sahl, 2006).	
	Advantages over antibiotics	Possible disadvantages
	<p>Selective antibacterial, antifungal, antiviral, anticancer, antiplasmodial, antiprotozoal, insecticidal, and spermicidal agents with no activities against host cells (Zhang &amp; Gallo, 2016). Not prone to resistance development</p>	<p>High toxicity (Zhang &amp; Gallo, 2016). Difficult industrial scalability (Fosgerau &amp; Hoffmann, 2015). Expensive large-scale production. Susceptible to proteolysis (Fosgerau &amp; Hoffmann, 2015).</p>
Strategy	Based on	
Bacteriocins	<p>Bacteriocins are ribosomal synthesized small AMPs produced by many bacterial species as a self-defense mechanism against other bacteria within the population to increase their probability of survival (Rea et al., 2011). Different modes of action are exhibited by bacteriocins, including suppression of peptidoglycan production and forming pores in cell membranes (Cotter et al., 2013).</p>	
	Advantages over antibiotics	Possible disadvantages
	<p>Selective against some bacteria (Rea et al., 2011). Active against drug-resistant pathogens. Have high tolerance towards harsh conditions, such as heat and UV exposure. Thus, Bacteriocins have many applications in the food industry (Cotter et al., 2013).</p>	<p>Bacterial pathogens can develop resistance to bacteriocins (Ghosh et al., 2019). Expensive large-scale production (Ghosh et al., 2019). Susceptible to proteolysis (Ghosh et al., 2019).</p>

<b>(2) Synthetically Generated Strategies</b>		
Strategy	Based on	
Synthetic mimics of antimicrobial peptides (SMAMPs)	SMAMPs are simple, rapidly bactericidal molecules, that are synthesized to maintain the properties of antimicrobial peptides while overcoming their disadvantages (Ghosh et al., 2019).	
	Advantages over antibiotics	Possible disadvantages
	Can turn back multi-drug resistant bacteria re-sensitive to antibiotics (Ghosh et al., 2019). Ease of synthesis (Ghosh et al., 2019). Broad-spectrum activity (Ghosh et al., 2019).	Toxicity (Konai et al., 2018). Challenges in administration (Ghosh et al., 2019). Bacterial pathogens can develop resistance to SMAMPs (Ghosh et al., 2019).
Strategy	Based on	
Antibacterial oligonucleotides	Oligonucleotides have been used to treat many infectious diseases, where antisense oligonucleotides with sequences complementary to resistance-causing genes have been used to silence these genes, in what is known as gene silencing therapy. It is an alternative strategy against multidrug-resistant bacteria (Ayhan et al., 2016).	
Strategy	Based on	
Innate defense regulatory peptides (IDRPs)	IDRPs are antiendotoxin and immunomodulatory peptides with no direct antibacterial activity. A promising candidate to conventional antibiotics with proven activity against bacterial and malarial infections (Hilchie et al., 2013).	
	Advantages over antibiotics	Possible disadvantages
	Modulate the immunesystem (Ghosh et al., 2019). No resistance development (Ghosh et al., 2019).	Expensive production (Ghosh et al., 2019). Susceptible to proteolysis (Ghosh et al., 2019).
Strategy	Based on	

Inhibitors of Bacterial Virulence	<p>The extracellular and cell-surface molecules that the bacterial pathogen produces to establish the infection are known as virulence factors. By targeting these virulence factors, the interaction between the bacterium and its host can be disrupted, thereby impeding the development of resistance in the bacterium. Consequently, this approach facilitates the elimination of the bacterium by the host immune system (Ghosh et al., 2019).</p>	
<b>(3) Biotechnology-Based Approaches</b>		
Strategy	Based on	
Genetically Modified Bacteriophages	<p>Genetically modified phages have been considered recently as an alternative strategy (Braff et al., 2016). With many modes of action, such as producing bacterial-biofilm-degrading enzymes (Lu &amp; Collins, 2007). This particular approach has the potential to reverse the resistance of microorganisms, such as <i>P. aeruginosa</i>, rendering them susceptible to standard antibiotics (Libis et al., 2014; Lu &amp; Collins, 2009).</p>	
Strategy	Based on	
Lysins (Endolysins, Exolysins, and Autolysins)	<p>One potential therapeutic approach that exhibits direct antibacterial activity is the utilization of either: 1) <b>Endolysins</b> generated by phages that facilitate the degradation of the bacterial cell wall (Nelson et al., 2012; Schmelcher et al., 2012), 2) <b>Exolysins</b> produced by bacteria to eliminate cells belonging to different strains or species or, 3) <b>Autolysins</b> which are synthesized during cellular growth and division (Basso et al., 2017).</p>	
	Advantages over antibiotics	Possible disadvantages
	<p>Vulnerable to genetic manipulation.                  Selective towards particular microorganisms (Ghosh et al., 2019).                  Not prone to resistance development.</p>	<p>Production challenges (Ghosh et al., 2019).</p>
Strategy	Based on	
CRISPR-Cas <sub>9</sub>	<p>The CRISPR (clustered, regularly interspaced, short palindromic repeats)-Cas9 systems comprise essential parts in the bacterial immune system, where 20 nt tiny RNA serves as a template for Cas9 to eliminate foreign genetic material at precise locations (Bikard &amp; Barrangou, 2017). The CRISPR-Cas9 system was</p>	

	delivered using bacteriophages to specifically target multidrug-resistant bacteria, rendering them susceptible (M. Goren et al., 2017).	
	Advantages over antibiotics	Possible disadvantages
	Specificity to certain strains (M. Goren et al., 2017; M. G. Goren et al., 2015).	Expensive production (Ghosh et al., 2019; Wang et al., 2016). Toxicity (Ghosh et al., 2019).

Some of these approaches show encouraging progress, yet many of them face many limitations usually related to; high production costs, toxicity, instability, and being strain- or species-specific rather than having the broad-spectrum effects of traditional antibiotics. For this reason, multiple therapeutics might be needed concurrently to treat wound infections.

### **5. Photodynamic therapy as a promising phototherapeutic approach for wound management**

One interesting option is a simple, non-invasive, and inexpensive phototherapeutic method that causes no significant tissue damage known as laser-based antimicrobial photodynamic therapy (aPDT) (Dai et al., 2009; Ghorbani et al., 2018; Gois et al., 2010). Moreover, it is unlikely that bacteria will become resistant to aPDT (Gupta et al., 2018; Schastak et al., 2010). Phototherapy has many approaches based on the interaction between light photons and different tissues, one of which is laser-based aPDT, a type of photochemotherapy initially developed for cancer treatment (Mang, 2004; Mitton & Ackroyd, 2008).

Lasers use the physical phenomena of stimulated emission to generate a light beam that is monochromatic, coherent, and collimated, characterized by a small bandwidth and low divergence (MAIMAN, 1960). Lasers exhibit superior efficiency for aPDT in comparison to alternative light sources due to these three characteristics (Hode, 2005). The therapeutic effects of laser-based phototherapy depend on the irradiation parameters of the laser output including fluence, intensity, mode of action either continuous wave or pulsed, exposure duration, and most importantly, its wavelength. The selection of a laser type for a specific



application is determined by these criteria (Medrado et al., 2003; Samaneh et al., 2015).

Laser-based aPDT refers to the process of eradicating specific pathogens by the photochemical stimulation of naturally occurring Photosensitizers (PS) molecules. These molecules can become active after being exposed to a specific wavelength of light (Sternberg et al., 1998), which then triggers photochemical or photophysical processes within the microbial cell. The therapeutic efficacy of photodynamic therapy (PDT) relies on the in situ generation of cytotoxic reactive oxygen species (ROS), which cause oxidative damage to various cellular components, ultimately resulting in cell death (de Annunzio et al., 2018; Ravanat et al., 2000; Stark, 2005). ROS are generated via two photochemical processes that occur simultaneously (Hamblin & Hasan, 2004). The first process involves electron transfer (type I), resulting in the formation of oxygen, peroxide, or hydroxide radicals. The second process involves energy transfer reactions (type II), leading to the production of singlet oxygen ( $^1O_2$ ) (Dolmans et al., 2003). These processes depend on the interaction between laser radiation and matter, which converts light energy into chemical energy (photochemical interaction) (Gois et al., 2010; Hanakova et al., 2014).

Femtosecond laser pulses have the unique ability to deposit energy into a microscopic volume on a very short time scale within a single laser pulse without affecting the surrounding surface of the tissue (El-Khordagui et al., 2017; RASHIDI et al., 2015). Both the pulse width and the repetition rate determine the heat accumulation in biological samples. If the heating duration is short, then the thermal energy can diffuse away more rapidly. And, if the duration between two pulses is long enough for the heat produced by the previous pulse to decay away, there will be no accumulation of heat (C. D. Clark et al., 2013). Femtosecond laser-based aPDT is one of the most promising emerging technologies for chronic wound management and infection as demonstrated in Table 3. Regrettably, an important challenge lies in determining the optimal wavelength, light dose, and fluence to effectively restrict bacterial growth while minimizing harm to the adjacent host tissue

**Table 3:** Investigating the bactericidal efficacy of femtosecond laser irradiation against different bacterial pathogens.

Light Source	Tested Organisms	External Photosensitizers	Irradiation Wavelength	Irradiation Parameters	Ref.
INSPIRE HF100 laser system (Spectra Physics) which was pumped by a mode-locked femtosecond Ti: sapphire laser MAI TAI HP (Spectra Physics)	- <i>Pseudomonas aeruginosa</i> - <i>Staphylococcus aureus</i>	-	370, 380, 390, 400, 410, 420, 480, 495, 700, and 800 nm	Power density = 0.047, 0.0377, 0.0354, and 0.0348 W/cm <sup>2</sup> Irradiation time = 1, 3, 5, 5.5, 7, 9, 10, 10.5, 15, and 20 min	(Ahmed et al., 2020)
	- <i>Staphylococcus aureus</i>	-	380, 390, 400, 410, and 420 nm	Average Power = 75, 150, and 210 mW Irradiation time = 5, 10 and 15 min. Energy density = 79.6, 159, 223, 239, 318, 420, 477, and 630 J/cm <sup>2</sup> .	(Ahmed et al., 2021)
	- <i>Pseudomonas aeruginosa</i> - <i>Listeria monocytogenes</i> - Methicillin-resistant <i>Staphylococcus aureus</i> (MRSA)	-	400 nm	Power = 50 mW Irradiation time = 15 min. Energy density = 160 J/cm <sup>2</sup>	(El-Gendy et al., 2021)
	- <i>Pseudomonas aeruginosa</i> - Methicillin-resistant <i>Staphylococcus aureus</i> (MRSA)	-	400 nm	Power = 50 mW Irradiation time = 15 min. Energy density = 160 J/cm <sup>2</sup>	(El-Gendy, Nawaf et al.)

	<ul style="list-style-type: none"> <li>- <i>Candida albicans</i></li> <li>- <i>Escherichia coli</i></li> <li>- <i>Enterococcus faecalis</i></li> <li>- <i>Listeria monocytogenes</i></li> <li>- Methicillin-resistant <i>Staphylococcus aureus</i> (MRSA)</li> </ul>	-	400 nm	<p>Power = 50 mW Irradiation time = 15 min. Energy density = 160 J/cm<sup>2</sup></p>	(El-Gendy, Obaid, et al., 2022)
--	--	---	--------	---	---------------------------------

## 6. Conclusion and future perspectives

Bacterial infections pose a significant health burden globally and with the increasing ineffectiveness of conventional treatments alternative approaches are urgently needed. Laser-based aPDT has emerged as a promising alternative for infection management. Additional research is required to investigate the potential clinical uses of various laser systems against specific microbial populations in tissues, such as microbial biofilm and the microbial expression of distinct virulence factors.

**Funding** This research received no external funding.

**Data availability** No datasets were generated or analysed during the current study.

**Declarations**

**Conflict of interest** The authors declare no conflict of interest.

**Competing interests** The authors declare no competing interests.

## References

- Abedon, S. T. (2015). Phage therapy of pulmonary infections. *Bacteriophage*, 5(1), e1020260. <https://doi.org/10.1080/21597081.2015.1020260>
- Abedon, S. T., Kuhl, S. J., Blasdel, B. G., & Kutter, E. M. (2011). Phage treatment of human infections. *Bacteriophage*, 1(2), 66–85. <https://doi.org/10.4161/bact.1.2.15845>
- Agren, M. S., Eaglstein, W. H., Ferguson, M. W., Harding, K. G., Moore, K., Saarialho-Kere, U. K., & Schultz, G. S. (2000). Causes and effects of the chronic inflammation in venous leg ulcers. *Acta Dermato-Venereologica. Supplementum*, 210, 3–17. <http://www.ncbi.nlm.nih.gov/pubmed/10884942>
- Ahmed, E., El-Gendy, A. O., Hamblin, M. R., & Mohamed, T. (2021). The effect of femtosecond laser irradiation on the growth kinetics of *Staphylococcus aureus*: An in vitro study. *Journal of Photochemistry and Photobiology B: Biology*, 221, 112240. <https://doi.org/10.1016/j.jphotobiol.2021.112240>
- Ahmed, E., El-Gendy, A. O., Moniem Radi, N. A., & Mohamed, T. (2020). The bactericidal efficacy of femtosecond laser-based therapy on the most common infectious bacterial pathogens in chronic wounds: an in vitro study. *Lasers in Medical Science*, ii. <https://doi.org/10.1007/s10103-020-03104-0>
- Amini, A., Soleimani, H., Rezaei, F., Ghoreishi, S. K., Chien, S., & Bayat, M. (2021). The Combined Effect of Photobiomodulation and Curcumin on Acute Skin Wound Healing in Rats. *Journal of Lasers in Medical Sciences*, 12(SE-Original Article), e9. <https://journals.sbmu.ac.ir/jlms/article/view/29555>
- Atkin, L., & Ousey, K. (2016). Wound bed preparation: A novel approach using HydroTherapy. *British Journal of Community Nursing*, 21(Sup12), S23–S28. <https://doi.org/10.12968/bjcn.2016.21.Sup12.S23>
- AYELLO, E. A., DOWSETT, C., SCHULTZ, G. S., SIBBALD, R. G., FALANGA, V., HARDING, K., ROMANELLI, M., STACEY, M., TÉOT, L., & VANSCHIEDT, W. (2004). TIME heals all wounds. *Nursing*, 34(4), 36–42. <https://doi.org/10.1097/00152193-200404000-00040>

- Ayhan, D. H., Tamer, Y. T., Akbar, M., Bailey, S. M., Wong, M., Daly, S. M., Greenberg, D. E., & Toprak, E. (2016). Sequence-Specific Targeting of Bacterial Resistance Genes Increases Antibiotic Efficacy. *PLOS Biology*, 14(9), e1002552. <https://doi.org/10.1371/journal.pbio.1002552>
- Barchitta, M., Maugeri, A., Favara, G., Magnano San Lio, R., Evola, G., Agodi, A., & Basile, G. (2019). Nutrition and Wound Healing: An Overview Focusing on the Beneficial Effects of Curcumin. *International Journal of Molecular Sciences*, 20(5), 1119. <https://doi.org/10.3390/ijms20051119>
- Barsoumian, A. E., Mende, K., Sanchez, C. J., Beckius, M. L., Wenke, J. C., Murray, C. K., & Akers, K. S. (2015). Clinical infectious outcomes associated with biofilm-related bacterial infections: a retrospective chart review. *BMC Infectious Diseases*, 15, 223. <https://doi.org/10.1186/s12879-015-0972-2>
- Basso, P., Ragno, M., Elsen, S., Reboud, E., Golovkine, G., Bouillot, S., Huber, P., Lory, S., Faudry, E., & Attrée, I. (2017). *Pseudomonas aeruginosa* Pore-Forming Exolysin and Type IV Pili Cooperate To Induce Host Cell Lysis. *MBio*, 8(1). <https://doi.org/10.1128/mBio.02250-16>
- Bikard, D., & Barrangou, R. (2017). Using CRISPR-Cas systems as antimicrobials. *Current Opinion in Microbiology*, 37, 155–160. <https://doi.org/10.1016/j.mib.2017.08.005>
- Boateng, J. S., Matthews, K. H., Stevens, H. N. E., & Eccleston, G. M. (2008). Wound Healing Dressings and Drug Delivery Systems: A Review. *Journal of Pharmaceutical Sciences*, 97(8), 2892–2923. <https://doi.org/10.1002/jps.21210>
- Boucher, H. W., Talbot, G. H., Bradley, J. S., Edwards, J. E., Gilbert, D., Rice, L. B., Scheld, M., Spellberg, B., & Bartlett, J. (2009). Bad Bugs, No Drugs: No ESKAPE! An Update from the Infectious Diseases Society of America. *Clinical Infectious Diseases*, 48(1), 1–12. <https://doi.org/10.1086/595011>
- Bowler, P. G., Duerden, B. I., & Armstrong, D. G. (2001). Wound microbiology and associated approaches to wound management. *Clinical Microbiology Reviews*, 14(2), 244–269. <https://doi.org/10.1128/CMR.14.2.244-269.2001>

- Braff, D., Shis, D., & Collins, J. J. (2016). Synthetic biology platform technologies for antimicrobial applications. *Advanced Drug Delivery Reviews*, 105, 35–43. <https://doi.org/10.1016/j.addr.2016.04.006>
- Brogden, K. A. (2005). Antimicrobial peptides: pore formers or metabolic inhibitors in bacteria? *Nature Reviews Microbiology*, 3(3), 238–250. <https://doi.org/10.1038/nrmicro1098>
- Cardona, A. F., & Wilson, S. E. (2015). Skin and Soft-Tissue Infections: A Critical Review and the Role of Telavancin in Their Treatment. *Clinical Infectious Diseases*, 61(suppl 2), S69–S78. <https://doi.org/10.1093/cid/civ528>
- Chang, R. Y. K., Morales, S., Okamoto, Y., & Chan, H.-K. (2020). Topical application of bacteriophages for treatment of wound infections. *Translational Research*, 220, 153–166. <https://doi.org/10.1016/j.trsl.2020.03.010>
- Clark, C. D., Marshall, W. J., & Thomas, R. J. (2013). Theoretical analysis of multiple-pulse thermal damage thresholds of the retina. *Journal of Laser Applications*, 25(1), 012005. <https://doi.org/10.2351/1.4788755>
- Clark, J. R., & March, J. B. (2006). Bacteriophages and biotechnology: vaccines, gene therapy and antibacterials. *Trends in Biotechnology*, 24(5), 212–218. <https://doi.org/10.1016/j.tibtech.2006.03.003>
- Coggan, K. A., & Wolfgang, M. C. (2012). Global regulatory pathways and cross-talk control pseudomonas aeruginosa environmental lifestyle and virulence phenotype. *Current Issues in Molecular Biology*, 14(2), 47–70. <http://www.ncbi.nlm.nih.gov/pubmed/22354680>
- Cooper, B., & Bachoo, P. (2018). Extracorporeal shock wave therapy for the healing and management of venous leg ulcers. *Cochrane Database of Systematic Reviews*, 2018(6). <https://doi.org/10.1002/14651858.CD011842.pub2>
- Cotter, P. D., Ross, R. P., & Hill, C. (2013). Bacteriocins — a viable alternative to antibiotics? *Nature Reviews Microbiology*, 11(2), 95–105. <https://doi.org/10.1038/nrmicro2937>
- Cullum, N., Nelson, E., Fletcher, A., & Sheldon, T. (2001). Compression for venous leg ulcers. In N. Cullum (Ed.), *The Cochrane Database of*

Systematic Reviews. John Wiley & Sons, Ltd.  
<https://doi.org/10.1002/14651858.CD000265>

- Dai, T., Huang, Y.-Y., & Hamblin, M. R. (2009). Photodynamic therapy for localized infections—State of the art. *Photodiagnosis and Photodynamic Therapy*, 6(3–4), 170–188. <https://doi.org/10.1016/j.pdpdt.2009.10.008>
- de Annunzio, S. R., de Freitas, L. M., Blanco, A. L., da Costa, M. M., Carmona-Vargas, C. C., de Oliveira, K. T., & Fontana, C. R. (2018). Susceptibility of *Enterococcus faecalis* and *Propionibacterium acnes* to antimicrobial photodynamic therapy. *Journal of Photochemistry and Photobiology B: Biology*, 178, 545–550. <https://doi.org/10.1016/j.jphotobiol.2017.11.035>
- Demidova-Rice, T. N., Durham, J. T., & Herman, I. M. (2012). Wound Healing Angiogenesis: Innovations and Challenges in Acute and Chronic Wound Healing. *Advances in Wound Care*, 1(1), 17–22. <https://doi.org/10.1089/wound.2011.0308>
- Dolmans, D. E. J. G. J., Fukumura, D., & Jain, R. K. (2003). Photodynamic therapy for cancer. *Nature Reviews Cancer*, 3(5), 380–387. <https://doi.org/10.1038/nrc1071>
- Dowd, S. E., Sun, Y., Secor, P. R., Rhoads, D. D., Wolcott, B. M., James, G. A., & Wolcott, R. D. (2008). Survey of bacterial diversity in chronic wounds using pyrosequencing, DGGE, and full ribosome shotgun sequencing. *BMC Microbiology*, 8, 43. <https://doi.org/10.1186/1471-2180-8-43>
- El-Gendy, A. O., Nawaf, K. T., Ahmed, E., Samir, A., Hamblin, M. R., Hassan, M., & Mohamed, T. (2022). Preparation of zinc oxide nanoparticles using laser-ablation technique: Retinal epithelial cell (ARPE-19) biocompatibility and antimicrobial activity when activated with femtosecond laser. *Journal of Photochemistry and Photobiology B: Biology*, 234, 112540. <https://doi.org/10.1016/j.jphotobiol.2022.112540>
- El-Gendy, A. O., Obaid, Y., Ahmed, E., Enwemeka, C. S., Hassan, M., & Mohamed, T. (2022). The Antimicrobial Effect of Gold Quantum Dots and Femtosecond Laser Irradiation on the Growth Kinetics of Common Infectious Eye Pathogens: An In Vitro Study. *Nanomaterials*, 12(21), 3757. <https://doi.org/10.3390/nano12213757>

- El-Gendy, A. O., Samir, A., Ahmed, E., Enwemeka, C. S., & Mohamed, T. (2021). The antimicrobial effect of 400 nm femtosecond laser and silver nanoparticles on gram-positive and gram-negative bacteria. *Journal of Photochemistry and Photobiology B: Biology*, 223, 112300. <https://doi.org/10.1016/j.jphotobiol.2021.112300>
- El-Khordagui, L., El-Sayed, N., Galal, S., El-Gowell, H., Omar, H., & Mohamed, M. (2017). Photosensitizer-eluting nanofibers for enhanced photodynamic therapy of wounds: A preclinical study in immunocompromized rats. *International Journal of Pharmaceutics*, 520(1–2), 139–148. <https://doi.org/10.1016/j.ijpharm.2017.02.004>
- Enoch, S., & Leaper, D. J. (2008). Basic science of wound healing. *Surgery (Oxford)*, 26(2), 31–37. <https://doi.org/10.1016/j.mpsur.2007.11.005>
- Fang, Y., Liu, T., Zou, Q., Zhao, Y., & Wu, F. (2016). Water-soluble benzylidene cyclopentanone based photosensitizers for in vitro and in vivo antimicrobial photodynamic therapy. *Scientific Reports*, 6(April), 1–9. <https://doi.org/10.1038/srep28357>
- Fosgerau, K., & Hoffmann, T. (2015). Peptide therapeutics: current status and future directions. *Drug Discovery Today*, 20(1), 122–128. <https://doi.org/10.1016/j.drudis.2014.10.003>
- Freidouni, M., Nejati, H., Salimi, M., Bayat, M., Amini, A., Noruzian, M., Ali Asgharie, M., & Rezaian, M. (2015). Evaluating Glucocorticoid Administration on Biomechanical Properties of Rats' Tibial Diaphysis. *Iranian Red Crescent Medical Journal*, 17(3). <https://doi.org/10.5812/ircmj.19389>
- Friedman, N. D., Temkin, E., & Carmeli, Y. (2016). The negative impact of antibiotic resistance. *Clinical Microbiology and Infection*, 22(5), 416–422. <https://doi.org/10.1016/j.cmi.2015.12.002>
- Frykberg, R. G., & Banks, J. (2015). Challenges in the Treatment of Chronic Wounds. *Advances in Wound Care*, 4(9), 560–582. <https://doi.org/10.1089/wound.2015.0635>
- Gallo, R. L., & Hooper, L. V. (2012). Epithelial antimicrobial defence of the skin and intestine. *Nature Reviews. Immunology*, 12(7), 503–516. <https://doi.org/10.1038/nri3228>



- Gao, X., & Xing, D. (2009). Molecular mechanisms of cell proliferation induced by low power laser irradiation. *Journal of Biomedical Science*, 16(1), 4. <https://doi.org/10.1186/1423-0127-16-4>
- Ghorbani, J., Rahban, D., Aghamiri, S., Teymouri, A., & Bahador, A. (2018). Photosensitizers in antibacterial photodynamic therapy: an overview. *Laser Therapy*, 27(4), 293–302. [https://doi.org/10.5978/islsm.27\\_18-RA-01](https://doi.org/10.5978/islsm.27_18-RA-01)
- Ghosh, C., Sarkar, P., Issa, R., & Haldar, J. (2019). Alternatives to Conventional Antibiotics in the Era of Antimicrobial Resistance. *Trends in Microbiology*, 27(4), 323–338. <https://doi.org/10.1016/j.tim.2018.12.010>
- Glass, G. E., Murphy, G. F., Esmaeili, A., Lai, L.-M., & Nanchahal, J. (2014). Systematic review of molecular mechanism of action of negative-pressure wound therapy. *British Journal of Surgery*, 101(13), 1627–1636. <https://doi.org/10.1002/bjs.9636>
- Gois, M. M., Kurachi, C., Santana, E. J. B., Mima, E. G. O., Spolidório, D. M. P., Pelino, J. E. P., & Salvador Bagnato, V. (2010). Susceptibility of *Staphylococcus aureus* to porphyrin-mediated photodynamic antimicrobial chemotherapy: an in vitro study. *Lasers in Medical Science*, 25(3), 391–395. <https://doi.org/10.1007/s10103-009-0705-0>
- Goren, M. G., Yosef, I., & Qimron, U. (2015). Programming Bacteriophages by Swapping Their Specificity Determinants. *Trends in Microbiology*, 23(12), 744–746. <https://doi.org/10.1016/j.tim.2015.10.006>
- Goren, M., Yosef, I., & Qimron, U. (2017). Sensitizing pathogens to antibiotics using the CRISPR-Cas system. *Drug Resistance Updates : Reviews and Commentaries in Antimicrobial and Anticancer Chemotherapy*, 30, 1–6. <https://doi.org/10.1016/j.drup.2016.11.001>
- Gravitz, L. (2012). Turning a new phage. *Nature Medicine*, 18(9), 1318–1320. <https://doi.org/10.1038/nm0912-1318>
- Guo, S., & Dipietro, L. A. A. (2010). Factors Affecting Wound Healing. *Journal of Dental Research*, 89(3), 219–229. <https://doi.org/10.1177/0022034509359125>
- Gupta, A., Keshri, G. K., & Yadav, A. (2018). Non-thermal Therapeutic Applications of Light. *Proceedings of the National Academy of Sciences*,

- India Section A: Physical Sciences, 88(3), 473–478.  
<https://doi.org/10.1007/s40010-018-0521-7>
- Hamblin, M. R., & Hasan, T. (2004). Photodynamic therapy: a new antimicrobial approach to infectious disease? *Photochemical & Photobiological Sciences : Official Journal of the European Photochemistry Association and the European Society for Photobiology*, 3(5), 436–450.  
<https://doi.org/10.1039/b311900a>
- Hanakova, A., Bogdanova, K., Tomankova, K., Pizova, K., Malohlava, J., Binder, S., Bajgar, R., Langova, K., Kolar, M., Mosinger, J., & Kolarova, H. (2014). The application of antimicrobial photodynamic therapy on *S. aureus* and *E. coli* using porphyrin photosensitizers bound to cyclodextrin. *Microbiological Research*, 169(2–3), 163–170.  
<https://doi.org/10.1016/j.micres.2013.07.005>
- Hancock, R. E. W., & Sahl, H.-G. (2006). Antimicrobial and host-defense peptides as new anti-infective therapeutic strategies. *Nature Biotechnology*, 24(12), 1551–1557. <https://doi.org/10.1038/nbt1267>
- Hilchie, A. L., Wuerth, K., & Hancock, R. E. W. (2013). Immune modulation by multifaceted cationic host defense (antimicrobial) peptides. *Nature Chemical Biology*, 9(12), 761–768.  
<https://doi.org/10.1038/nchembio.1393>
- Hode, L. (2005). The Importance of the Coherency. *Photomedicine and Laser Surgery*, 23(4), 431–434. <https://doi.org/10.1089/pho.2005.23.431>
- Hosseinkhani, A., Falahatzadeh, M., Raoofi, E., & Zarshenas, M. M. (2017). An Evidence-Based Review on Wound Healing Herbal Remedies From Reports of Traditional Persian Medicine. *Journal of Evidence-Based Complementary & Alternative Medicine*, 22(2), 334–343.  
<https://doi.org/10.1177/2156587216654773>
- Jackson, N., Czaplewski, L., & Piddock, L. J. V. (2018). Discovery and development of new antibacterial drugs: learning from experience? *The Journal of Antimicrobial Chemotherapy*, 73(6), 1452–1459.  
<https://doi.org/10.1093/jac/dky019>
- Jones, M. E., Karlowsky, J. A., Draghi, D. C., Thornsberry, C., Sahm, D. F., & Nathwani, D. (2003). Epidemiology and antibiotic susceptibility of bacteria causing skin and soft tissue infections in the USA and Europe: a

- guide to appropriate antimicrobial therapy. *International Journal of Antimicrobial Agents*, 22(4), 406–419. [https://doi.org/10.1016/S0924-8579\(03\)00154-7](https://doi.org/10.1016/S0924-8579(03)00154-7)
- Jones, R. N. (2001). Resistance patterns among nosocomial pathogens: trends over the past few years. *Chest*, 119(2 Suppl), 397S-404S. [https://doi.org/10.1378/chest.119.2\\_suppl.397s](https://doi.org/10.1378/chest.119.2_suppl.397s)
- Kavros, S. J., & Coronado, R. (2018). Diagnostic and Therapeutic Ultrasound on Venous and Arterial Ulcers: A Focused Review. *Advances in Skin & Wound Care*, 31(2), 55–65. <https://doi.org/10.1097/01.ASW.0000527967.10613.87>
- Khouri, C., Kotzki, S., Roustit, M., Blaise, S., Gueyffier, F., & Cracowski, J.-L. (2017). Hierarchical evaluation of electrical stimulation protocols for chronic wound healing: An effect size meta-analysis. *Wound Repair and Regeneration*, 25(5), 883–891. <https://doi.org/10.1111/wrr.12594>
- Kloth, L. C. (2014). Electrical Stimulation Technologies for Wound Healing. *Advances in Wound Care*, 3(2), 81–90. <https://doi.org/10.1089/wound.2013.0459>
- Kmietowicz, Z. (2017). Few novel antibiotics in the pipeline, WHO warns. *BMJ (Clinical Research Ed.)*, 358, j4339. <https://doi.org/10.1136/bmj.j4339>
- Kohanski, M. A., Dwyer, D. J., & Collins, J. J. (2010). How antibiotics kill bacteria: from targets to networks. *Nature Reviews Microbiology*, 8(6), 423–435. <https://doi.org/10.1038/nrmicro2333>
- Konai, M. M., Bhattacharjee, B., Ghosh, S., & Haldar, J. (2018). Recent Progress in Polymer Research to Tackle Infections and Antimicrobial Resistance. *Biomacromolecules*, 19(6), 1888–1917. <https://doi.org/10.1021/acs.biomac.8b00458>
- Kraus, C. N. (2008). Low hanging fruit in infectious disease drug development. *Current Opinion in Microbiology*, 11(5), 434–438. <https://doi.org/10.1016/j.mib.2008.09.009>
- Kuffler, D. P. (2016). Photobiomodulation in promoting wound healing: a review. *Regenerative Medicine*, 11(1), 107–122. <https://doi.org/10.2217/rme.15.82>

- Kwan, R. L.-C., Wong, W.-C., Yip, S.-L., Chan, K.-L., Zheng, Y.-P., & Cheing, G. L.-Y. (2015). Pulsed Electromagnetic Field Therapy Promotes Healing and Microcirculation of Chronic Diabetic Foot Ulcers. *Advances in Skin & Wound Care*, 28(5), 212–219. <https://doi.org/10.1097/01.ASW.0000462012.58911.53>
- Labrie, S. J., Samson, J. E., & Moineau, S. (2010). Bacteriophage resistance mechanisms. *Nature Reviews Microbiology*, 8(5), 317–327. <https://doi.org/10.1038/nrmicro2315>
- Lalezari, S., Lee, C. J., Borovikova, A. A., Banyard, D. A., Paydar, K. Z., Wirth, G. A., & Widgerow, A. D. (2017). Deconstructing negative pressure wound therapy. *International Wound Journal*, 14(4), 649–657. <https://doi.org/10.1111/iwj.12658>
- Lambrechts, S. A. G., Demidova, T. N., Aalders, M. C. G., Hasan, T., & Hamblin, M. R. (2005). Photodynamic therapy for *Staphylococcus aureus* infected burn wounds in mice. *Photochemical & Photobiological Sciences*, 4(7), 503. <https://doi.org/10.1039/b502125a>
- Las Heras, K., Igartua, M., Santos-Vizcaino, E., & Hernandez, R. M. (2020). Chronic wounds: Current status, available strategies and emerging therapeutic solutions. *Journal of Controlled Release*, 328, 532–550. <https://doi.org/10.1016/j.jconrel.2020.09.039>
- Libis, V. K., Bernheim, A. G., Basier, C., Jaramillo-Riveri, S., Deyell, M., Aghoghogbe, I., Atanaskovic, I., Bencherif, A. C., Benony, M., Koutsoubelis, N., Löchner, A. C., Marinkovic, Z. S., Zahra, S., Zegman, Y., Lindner, A. B., & Wintermute, E. H. (2014). Silencing of Antibiotic Resistance in *E. coli* with Engineered Phage Bearing Small Regulatory RNAs. *ACS Synthetic Biology*, 3(12), 1003–1006. <https://doi.org/10.1021/sb500033d>
- Livermore, D. M. (2007). Introduction: the challenge of multiresistance. *International Journal of Antimicrobial Agents*, 29 Suppl 3, S1-7. [https://doi.org/10.1016/S0924-8579\(07\)00158-6](https://doi.org/10.1016/S0924-8579(07)00158-6)
- Lu, T. K., & Collins, J. J. (2007). Dispersing biofilms with engineered enzymatic bacteriophage. *Proceedings of the National Academy of Sciences*, 104(27), 11197–11202. <https://doi.org/10.1073/pnas.0704624104>

- Lu, T. K., & Collins, J. J. (2009). Engineered bacteriophage targeting gene networks as adjuvants for antibiotic therapy. *Proceedings of the National Academy of Sciences*, 106(12), 4629–4634. <https://doi.org/10.1073/pnas.0800442106>
- MAIMAN, T. H. (1960). Stimulated Optical Radiation in Ruby. *Nature*, 187(4736), 493–494. <https://doi.org/10.1038/187493a0>
- Mang, T. S. (2004). Lasers and light sources for PDT: past, present and future. *Photodiagnosis and Photodynamic Therapy*, 1(1), 43–48. [https://doi.org/10.1016/S1572-1000\(04\)00012-2](https://doi.org/10.1016/S1572-1000(04)00012-2)
- Medrado, A. R. A. P., Pugliese, L. S., Reis, S. R. A., & Andrade, Z. A. (2003). Influence of low level laser therapy on wound healing and its biological action upon myofibroblasts. *Lasers in Surgery and Medicine*, 32(3), 239–244. <https://doi.org/10.1002/lsm.10126>
- Mitton, D., & Ackroyd, R. (2008). A brief overview of photodynamic therapy in Europe. *Photodiagnosis and Photodynamic Therapy*, 5(2), 103–111. <https://doi.org/10.1016/j.pdpdt.2008.04.004>
- Mizutani, K., Musya, Y., Wakae, K., Kobayashi, T., Tobe, M., Taira, K., & Harada, T. (2004). A Clinical Study on Serum Prostaglandin E 2 with Low-Level Laser Therapy. *Photomedicine and Laser Surgery*, 22(6), 537–539. <https://doi.org/10.1089/pho.2004.22.537>
- Mustoe, T. A., O’Shaughnessy, K., & Kloeters, O. (2006). Chronic wound pathogenesis and current treatment strategies: a unifying hypothesis. *Plastic and Reconstructive Surgery*, 117(7 Suppl), 35S–41S. <https://doi.org/10.1097/01.prs.0000225431.63010.1b>
- Nagori, B. P., & Solanki, R. (2011). Role of Medicinal Plants in Wound Healing. *Research Journal of Medicinal Plant*, 5(4), 392–405. <https://doi.org/10.3923/rjmp.2011.392.405>
- Nelson, D. C., Schmelcher, M., Rodriguez-Rubio, L., Klumpp, J., Pritchard, D. G., Dong, S., & Donovan, D. M. (2012). Endolysins as Antimicrobials (pp. 299–365). <https://doi.org/10.1016/B978-0-12-394438-2.00007-4>
- Nesi-Reis, V., Lera-Nonose, D. S. S. L., Oyama, J., Silva-Lalucci, M. P. P., Demarchi, I. G., Aristides, S. M. A., Teixeira, J. J. V., Silveira, T. G. V., & Lonardoni, M. V. C. (2018). Contribution of photodynamic therapy in

- wound healing: A systematic review. *Photodiagnosis and Photodynamic Therapy*, 21, 294–305. <https://doi.org/10.1016/j.pdpdt.2017.12.015>
- Nourian Dehkordi, A., Mirahmadi Babaheydari, F., Chehelgerdi, M., & Raeisi Dehkordi, S. (2019). Skin tissue engineering: wound healing based on stem-cell-based therapeutic strategies. *Stem Cell Research & Therapy*, 10(1), 111. <https://doi.org/10.1186/s13287-019-1212-2>
- Otero-Viñas, M., & Falanga, V. (2016). Mesenchymal Stem Cells in Chronic Wounds: The Spectrum from Basic to Advanced Therapy. *Advances in Wound Care*, 5(4), 149–163. <https://doi.org/10.1089/wound.2015.0627>
- Paterson, D. L. (2006). Resistance in gram-negative bacteria: enterobacteriaceae. *The American Journal of Medicine*, 119(6 Suppl 1), S20-8; discussion S62-70. <https://doi.org/10.1016/j.amjmed.2006.03.013>
- Pierce, G. F., & Mustoe, T. A. (1995). Pharmacologic enhancement of wound healing. *Annual Review of Medicine*, 46, 467–481. <https://doi.org/10.1146/annurev.med.46.1.467>
- Price, P., Bale, S., Crook, H., & Harding, K. G. H. (2000). The effect of a radiant heat dressing on pressure ulcers. *Journal of Wound Care*, 9(4), 201–205. <https://doi.org/10.12968/jowc.2000.9.4.25972>
- Qureshi, A. A., Ross, K. M., Ogawa, R., & Orgill, D. P. (2011). Shock Wave Therapy in Wound Healing. *Plastic and Reconstructive Surgery*, 128(6), 721e-727e. <https://doi.org/10.1097/PRS.0b013e318230c7d1>
- Rai, M., Kon, K., Gade, A., Ingle, A., Nagaonkar, D., Paralikar, P., & da Silva, S. S. (2016). Antibiotic Resistance. In *Antibiotic Resistance* (pp. 121–143). Elsevier. <https://doi.org/10.1016/B978-0-12-803642-6.00006-X>
- RASHIDI, S., YADOLLAHPOUR, A. Y., & MIRZAIYAN, M. (2015). Low Level Laser Therapy for the Treatment of Chronic Wound: Clinical Considerations. *Biomedical and Pharmacology Journal*, 8(2), 1121–1127. <https://doi.org/10.13005/bpj/866>
- Ravanat, J.-L., Di Mascio, P., Martinez, G. R., Medeiros, M. H. G., & Cadet, J. (2000). Singlet Oxygen Induces Oxidation of Cellular DNA. *Journal of Biological Chemistry*, 275(51), 40601–40604. <https://doi.org/10.1074/jbc.M006681200>
- Rea, M. C., Dobson, A., O’Sullivan, O., Crispie, F., Fouhy, F., Cotter, P. D., Shanahan, F., Kiely, B., Hill, C., & Ross, R. P. (2011). Effect of broad-

- and narrow-spectrum antimicrobials on *Clostridium difficile* and microbial diversity in a model of the distal colon. *Proceedings of the National Academy of Sciences*, 108(Supplement\_1), 4639–4644. <https://doi.org/10.1073/pnas.1001224107>
- Rice, L. B. (2006). Antimicrobial resistance in gram-positive bacteria. *The American Journal of Medicine*, 119(6 Suppl 1), S11-9; discussion S62-70. <https://doi.org/10.1016/j.amjmed.2006.03.012>
- Rice, L. B. (2008). Federal Funding for the Study of Antimicrobial Resistance in Nosocomial Pathogens: No ESKAPE. *The Journal of Infectious Diseases*, 197(8), 1079–1081. <https://doi.org/10.1086/533452>
- Samaneh, R., Ali, Y., Mostafa, J., Mahmud, N. A., & Zohre, R. (2015). Laser Therapy for Wound Healing: A Review of Current Techniques and Mechanisms of Action. *Biosciences Biotechnology Research Asia*, 12(SEMAR), 217–223. <https://doi.org/10.13005/bbra/1626>
- Schastak, S., Ziganshyna, S., Gitter, B., Wiedemann, P., & Claudepierre, T. (2010). Efficient Photodynamic Therapy against Gram-Positive and Gram-Negative Bacteria Using THPTS, a Cationic Photosensitizer Excited by Infrared Wavelength. *PLoS ONE*, 5(7), e11674. <https://doi.org/10.1371/journal.pone.0011674>
- Schmelcher, M., Donovan, D. M., & Loessner, M. J. (2012). Bacteriophage endolysins as novel antimicrobials. *Future Microbiology*, 7(10), 1147–1171. <https://doi.org/10.2217/fmb.12.97>
- Schultz, G. S., Sibbald, R. G., Falanga, V., Ayello, E. A., Dowsett, C., Harding, K., Romanelli, M., Stacey, M. C., Teot, L., & Vanscheidt, W. (2003). Wound bed preparation: a systematic approach to wound management. *Wound Repair and Regeneration: Official Publication of the Wound Healing Society [and] the European Tissue Repair Society*, 11 Suppl 1, S1-28. <https://doi.org/10.1046/j.1524-475x.11.s2.1.x>
- Sefati, N., Abbaszadeh, H.-A., Fadaei Fathabady, F., Abdollahifar, M.-A., Amini, A., Noori-Zadeh, A., Darabi, S., & Norouzian, M. (2018). Therapeutic Effects of Laser on Partial Osteotomy in the Rat Model of Hypothyroidism. *Journal of Lasers in Medical Sciences*, 9(2), 121–127. <https://doi.org/10.15171/jlms.2018.23>

- Segev-Zarko, L., Saar-Dover, R., Brumfeld, V., Mangoni, M. L., & Shai, Y. (2015). Mechanisms of biofilm inhibition and degradation by antimicrobial peptides. *Biochemical Journal*, 468(2), 259–270. <https://doi.org/10.1042/BJ20141251>
- Shpichka, A., Butnaru, D., Bezrukov, E. A., Sukhanov, R. B., Atala, A., Burdukovskii, V., Zhang, Y., & Timashev, P. (2019). Skin tissue regeneration for burn injury. *Stem Cell Research & Therapy*, 10(1), 94. <https://doi.org/10.1186/s13287-019-1203-3>
- Shrestha, G., Raphael, J., Leavitt, S. D., & St. Clair, L. L. (2014). In vitro evaluation of the antibacterial activity of extracts from 34 species of North American lichens. *Pharmaceutical Biology*, 52(10), 1262–1266. <https://doi.org/10.3109/13880209.2014.889175>
- Sibbald, R. G., Williamson, D., Orsted, H. L., Campbell, K., Keast, D., Krasner, D., & Sibbald, D. (2000). Preparing the wound bed--debridement, bacterial balance, and moisture balance. *Ostomy/Wound Management*, 46(11), 14–22, 24–28, 30–35; quiz 36–37. <http://www.ncbi.nlm.nih.gov/pubmed/11889735>
- Singer, A. J., & Clark, R. A. (1999). Cutaneous wound healing. *The New England Journal of Medicine*, 341(10), 738–746. <https://doi.org/10.1056/NEJM199909023411006>
- Spellberg, B., Guidos, R., Gilbert, D., Bradley, J., Boucher, H. W., Scheld, W. M., Bartlett, J. G., Edwards, J., & Infectious Diseases Society of America. (2008). The epidemic of antibiotic-resistant infections: a call to action for the medical community from the Infectious Diseases Society of America. *Clinical Infectious Diseases : An Official Publication of the Infectious Diseases Society of America*, 46(2), 155–164. <https://doi.org/10.1086/524891>
- Sprockett, D. D., Ammons, C. G., & Tuttle, M. S. (2015). Use of 16S rRNA sequencing and quantitative PCR to correlate venous leg ulcer bacterial bioburden dynamics with wound expansion, antibiotic therapy, and healing. *Wound Repair and Regeneration : Official Publication of the Wound Healing Society [and] the European Tissue Repair Society*, 23(5), 765–771. <https://doi.org/10.1111/wrr.12309>



- 
- Stark, G. (2005). Functional Consequences of Oxidative Membrane Damage. *Journal of Membrane Biology*, 205(1), 1–16. <https://doi.org/10.1007/s00232-005-0753-8>
- Steed, D. L. (2004). Debridement. *American Journal of Surgery*, 187(5A), 71S–74S. [https://doi.org/10.1016/S0002-9610\(03\)00307-6](https://doi.org/10.1016/S0002-9610(03)00307-6)
- Sternberg, E. D., Dolphin, D., & Brückner, C. (1998). Porphyrin-based photosensitizers for use in photodynamic therapy. *Tetrahedron*, 54(17), 4151–4202. [https://doi.org/10.1016/S0040-4020\(98\)00015-5](https://doi.org/10.1016/S0040-4020(98)00015-5)
- Thom, S. R. (2009). Oxidative stress is fundamental to hyperbaric oxygen therapy. *Journal of Applied Physiology*, 106(3), 988–995. <https://doi.org/10.1152/jappphysiol.91004.2008>
- Wang, H., La Russa, M., & Qi, L. S. (2016). CRISPR/Cas9 in Genome Editing and Beyond. *Annual Review of Biochemistry*, 85, 227–264. <https://doi.org/10.1146/annurev-biochem-060815-014607>
- Xue, M., & Jackson, C. J. (2015). Extracellular Matrix Reorganization During Wound Healing and Its Impact on Abnormal Scarring. *Advances in Wound Care*, 4(3), 119–136. <https://doi.org/10.1089/wound.2013.0485>
- Yildirimer, L., Thanh, N. T. K., & Seifalian, A. M. (2012). Skin regeneration scaffolds: a multimodal bottom-up approach. *Trends in Biotechnology*, 30(12), 638–648. <https://doi.org/10.1016/j.tibtech.2012.08.004>
- Zhang, L., & Gallo, R. L. (2016). Antimicrobial peptides. *Current Biology*, 26(1), R14–R19. <https://doi.org/10.1016/j.cub.2015.11.017>